Aspergillus Genomes Projects: Review, Resources, Uses and Future Applications

Ginsau, M.A
Department of Science Laboratory Technology,
Jigawa State Polytechnic,
Dutse, Nigeria.
Ginsau76@yahoo.com

***Abstract*-The genome sequence information has in recent times turned out to be available for setting Aspergillus species. The initial Aspergillus genome to be ordered is such of A. nidulans, with a three times exposure realized in 1998 by Cereon Genomics (Monsanto). This was prepared obtainable by confined community right to use in April 2001, as well as complete right to use approved in January 2003. A complete thirteen time's exposure of the A. nidulans genome, integrating the Cereon information, has from that time made available to public by the Whitehead Institute/MIT Center for Genome Research. No database holds each of the 14 available Aspergillus genomes; most are available from various resources. Aspergillus genomics is used in human diseases and biotechnology. The arrangements of Aspergillus genomes, together the right now available and the ones to be released sometime later will proceed to make the advancement of alternate levels of system-wide study. It recommended increasing the access of the genomes.**

Key words: **Aspergillus, genomes, projects, resources, uses, applications.**

I. INTRODUCTION

Initially depicted almost for the past three hundred years by Antonio Micheli, a cleric and botanist, *Aspergillus* received this title as a result of similarity of its asexual spore-makingfeature to the aspergillum, a device employed in churches to sprinkle blessed water in some liturgical activities.

Aspergillus is hence the title which depicts the asexual sequence of the fungus. Light of the fact that the phenotypic differing qualities of the sexual fruiting parts are bigger, ten diverse genuses portray the sexual sequence of *Aspergillus* species [23].

According to [4] the significance of the expression genomics have developed quickly and in no way it's utilized among fungi is unlike. Genomics can be defined as "the determination and use of the genome sequence of organisms to identify genes and non-coding, but potentially functional important, region of the genome". Post-genomics takes us more in utilizing the data to research function of genes inside in corporate cellular method subsequently giving insight interested in the molecular origin of the biology of living things; the data can likewise be applied to increase points of view on evolutionary associations among species. A variety of ways must be employed that incorporate the investigation of transcriptional procedures (transcriptomics), protein making and protein–protein relations (proteomics) and metabolism (metabolomics). It is essential to have the capacity to genetic copy particular parts of DNA for their useful description, toward positive and losing control the transcription of genes each incompletely or totally by gene knockout, to bring DNA interested in the genome regularly at precise genomic loci, to like the result of epigenetic impacts and to include screens to notice consequential changes. At the same time as a technical word, Genomics might have been newly instituted as well as useful in the direction of filamentous fungi; however, traditional genetics research with *Neurosporacrassa* and *Aspergilusnidulans* established the framework. Despite the fact that the periodical of fungal genome order started with the yeast *Saccharomyces cerevisiae*, the later periodical of the annotated *N. crassa* genome sequence and its complete pulling out, incorporating disclosures in relation to DNA methylation, have taken fungal genomics to a novel stage. The accessibility of a bigger genome in the company of the included intricacy emerging from the filamentous growth structure, as compared with yeast, has furnished novel chances for genome-wide research.

Genomic and post-genomic studies including *Aspergillus* species will supplement and put in to research with *N. crassa* by presenting a variety of species among differing natural attributes e.g. replica and practical, sexual and asexual, pathogenic and to high extent non-pathogenic organisms.

The genomic uprisings have been achieved through enhanced processes in favor of sequencing, producing libraries, and explanation. Following DNA sequencing, computerized explanation utilizes bioinformatics' gene discovery programmes to spot the protein coding portions of genomes. The program work most excellent after 'skilled' on the suitable genome. Computerized explanation can present an exceptional original outline of the gene substance and display of a genome. On the other hand, automatic annotation is famously inaccurate and subsequently must be subject to nonstop updates as more experimentally based data gets available. Following explanation, collected genes could be grouped as enzymes, receptors, and transcription factors.

IJRS

Explained genomes permit the analysis of genes fall starting the same predecessor over numerous diverse organisms. Occasionally this relative genomic methodology permits us togive supposed role to obscure anticipated genes. The synchronous announcement of three *Aspergillus* genome script in December 2005in *Nature* created *Aspergillus* as the foremost filamentous fungal genus in favor of relative genomic research [21]; [30]; [31].

II. ASPERGILLUS GENOMES

Genome sequence information has in recent time sturned out to be available for setting *Aspergillus* species. The initial *Aspergillus* genome to be ordered is such of *A. nidulans*, with a three times exposure realized in 1998 by Cereon Genomics (Monsanto). This was prepared obtainable by confined community right to use in April 2001, as well as complete right to use approved in January 2003. A complete thirteen time's exposure of the *A. nidulans* genome, integrating the Cereon information, has from that time made available to public by the Whitehead Institute/MIT Center for Genome Research [4].

Aspergillus is the extremely genome ordered prosperous fungal group, with the genomes from 14 species previously publically accessible [23]. Going beyond even the genome sequenced rich *Saccharomyces* and *Candida* yeasts[14].

Aspergillus is prone to keep possessing on to this difference since the Department of Energy Joint Genome Institute (JGI) of United State like component of its 2011 society ordering suggestion system, accepted a many-researcher offer and is at this time ordering the genomes of 8supplementary species, together with the coral pathogen *Asydowii* and the xerophile *Aspergillusgalaucus.*

III. ASPERGILLUS GENOME PROJECTS

In spite of the fact that genome sequencing projects for the *Aspergillus* species, *A.nidulans* and *Aspergillus oryzae* began as timely as 1998 and 1996, in that order, the genomics period of filamentous fungi is by and large settled to have begun in February 2001, when the Whitehead Institute Center for Genome Research published a draft form of the genome ordering of *N.crassa* [2]. [22] Published the finalized form of the genome. As a demonstration of the essentialness of the job, one hundred and seven page appraisal of the effect of the *N. crassa*genome ordering was published by [9].

Aspergillus genomics, therefore, began in January 2003, where complete free right to use was granted to a three times coverage *A. nidulans* genome sequence from Cereon genomics. Two months after, this sequence was published again by the Whitehead Institute/MIT Center for Genome Research with the expansion often times coverage ordering to produce an outline genome sequence with thirteen times coverage. Additionally, the Dutch organization DSM reported the genome ordering of an enzyme-producing strain of *Aspergillus Niger* (CBS 513.88) in a journalists release in December 2001, however, restricted right to use to collaborators. A periodical on the *A. nidulans* genome sequence was released in December 2005 together with periodicals on the genome ordering of *A. oryzae* and the opportunistic pathogen *Aspergillus fumigatus,* the principal cause behind persistent a spergillos is, the most recurrent fungal infection on the planet [31]; [30]; [21], whereas the genome order of *A. niger* CBS 513.88 was made available to the general society, in addition to the publication of [34].

Some *Aspergillus* genomes have been ordered in comparable with or throughout the landmark periodicals of these four genomes, a number of them by autonomous organizations, others like a component of the public effort.

Individual of these effort is the Fungal Genome Initiative (FGI) (http://www.genome. wi.mit.edu/seq/fgi/) which since 2002 has published white papers telling that fungi to be sequenced, incorporating the ten times sequencing of *A. nidulans* which was a component of the suggestion of the original white paper. Since that time, four white papers have been published signifying the sequencing of a sum of ninety nine fungal species, incorporating eight *Aspergilli*.

Table I: Facts of Some *Aspergillus* Species.

Aspergillus aculeatus	A plant pathogen likewise used on behalf of detection of novel extracellular enzymes, where exciting genes were duplicated from cDNA and expressed in *A. oryzae.*
Aspergillus carbonarius	Anochratox in maker which is associated in food decay, especially grapes. The organism is almost connected with *A. niger* and other dark *Aspergilli.*
Aspergillus clavatus	An allergenic and seldom pathogenic fungus which is fundamentally being ordered to add a supplementary comprehension of the two almost interrelated pathogenic *Aspergilli A. fumigates* and *Aspergillus terreus.*
Aspergillus (Neosartorya) fischeri	A non-pathogenic nearly linked to the pathogenic *A. fumigatus.*
Aspergillus flavus	Human being pathogen producing the quite cancer-causing aflatox in. It additionally generates finacial misfortunes because of the grain infection.
Aspergillus parasiticus	The maker of a flatoxin.
Aspergillus terreus	Used technologically to make the cholesterol, slowing down agent's pravastatin, simvastatin, and lovastatin and the organic acid I a conic acid, yet has additionally been associated with aspergillosis.

IJRS

[2] Also mentioned that as the genome sequence keeps on to increase, likewise the need for genome projects organizations.

Resources such as the Broad Institute *Aspergillus* comparative data base (http://www.broad.mit.edu/annotation/genome/ aspergillus_group/multihome.html), e-fungi (http://www.cs.man. ac.uk/~cornell/efungi/database.html), and the central *Aspergillus* data repository (CADRE) (http://www.cadre-genomes.org.uk/, intended for the similarity of genomic information, will debatably be of considerably more significance to investigators as the figure of the entire genome sequenced *Aspergilli* raises and allow further research dependent upon relative genomics. The initial4 point of reference genome documents have stimulated an extensive series of genome-motivated occupation, as might be deliberated in a later unique matter of researches in Mycology, (number fifty nine) on "*Aspergillus* systematics in the genomic era", and the current problem of Fungal genetics.

An outline of the position and possible of the genome motivated approach, the accompanying segments, will delineate research about genomics in *Aspergillus* species, wherever such has been published. According to perspective of fungi in [4] the later meaning of genomics was "the determination and use of the genome sequence of organisms to identify genes and non-coding, but potentially functionally important, regions of the genome". [2] concludes that this general idea will center on this sort of research, for example the publications on complete genome sequencing or genome surveys; in any case, a couple of cases of system-wide useful genomics or relative genomics will be incorporated to give prospects of *Aspergillus* genomics as follows:

Aspergillus Aculeatus Genomics: The genome sequence of *A. aculeatus* is yet in the ordering and explanation stage of the Joint Genome Institute (JGI) (Scott Baker, personal communications), and no investigation of the genomics or the prospects of these have been published at this point.

Aspergillus Carbonarius Genomics: Likewise with *A. aculeatus,* the genome sequence of *A. carbonarius* has not up till now been issued from the JGI (Scott Baker, personal communications). Genomic researches have been restricted to a relatively arning of the mitochondrial genome of several *A. carbonarius* strains based mainly on limitation mapping but not on genome sequencing.

Aspergillus Clavatus Genomics: The genome sequencing of *A. clavatus* strain NRRL 1 has in recent times been published by [20]together in the company of the genome sequences of *Neurosporafischeri* and a second strain of *A. fumigatus*, which is available from NCBI in its 3rdreport.

Study of this fungus is scanty contrasted with extra *Aspergilli,* and the inputs in light of the nucleotide order are constrained to the cloning of clavin. [40] Expected comparative genomics of *A. fumigatus, A. clavatus,* and *N. fischeri* will allow the detection of factors accountable for pathogen city in *A. fumigatus.* Pathogen city was for sure one of the things of research in the genome by [20] and will be talked about in particular in the part on *A. fumigatus* genomics.

Neurosporafischeri Genomics: The genome order of the 8 chromosomes of *N. fischeri* has been examined and compared with the genomes of *A. fumigatus*Af293, A1163 and *A. clavatus*NRRL 1 by [20].

Aspergillus Flavus Genomics: The preface remarks on the periodicals of the *A. flavus*genome have been published with the sequence data available by [40]. Although the article does not hold the genomic research discovered in the periodicals of *A. Niger, A. nidulans, A. fumigates,* and *A. oryzae,* a flatoxin biosynthesis and the aflatoxin gene group were specified as focuses of individual attention for upcoming research. This has, in fact, been investigated more, as will be depicted in the segment on transcriptomics.

Aspergillus Fumigates Genomics: The sequencing of *A. fumigatus*Af293 was started in 2001, and a remark was available by [16], telling the expectation of how the scheme might stimulate the research in anti-fungal drugs. The explanation and analysis of the 922 kb bulky area were published by [32], permitting the near the beginning recognition of components supposedly involved in pathogencity and anti-fungal functions. This research was discussed by the announcement of the study of the complete genome sequence by [31], where the genome was inspected for allergens and characteristics that could be involved in pathogenicity. These examinations anticipated 9 proteins to be allergens, and recognized 26 groups that may be involved in the generation of resultant metabolites.

Moreover, the copied of the medical isolate strain *A. fumigates*CEA10, *A. fumigatus*FGSC A1163, has been arranged by Merck & Co and was liable to a preparatory study by [40] and a relativestudy of *A. fumigatus*FGSCA1163 and *A. Fumigatus A*f293 by [35]. Out of the new periodical of the genome sequence of *A. fumigatus*FGSC A1163 along with the sequence of *A. clavatus*NRRL 1 and *N. fischeri*NRRL 181, the genome sequences were compared to recognize genomic components interesting to *A. fumigatus* FGSC A1163, and to *A. fumigatus*, accordingly giving possible focuses for additional investigation of the reason for pathogenicity.

The research incitement by the genome sequence of *A. fumigatus* Af293 covers numerous perspectives, as was described in a report by [36], focusing on an extensive variety of attributes and relative genomics to *A. oryzae* and *A. nidulans.* Single utilization of the genome order in the

IJRS

perspective of essential study was the recognition of mating-type- like genes in the asexual A. *fumigates* reviewed by [18].

The 2 mating breeds were at last affirmed by [33], accordingly recommending a dynamic sexual cycle. A study of the effect of the genome order of pathogenic fungi well as distinguishing proof of additional supposed pathogenic gene assembly has been advanced by [12].

Aspergillus Nidulans Genomics: As depicted in the prologue to *Aspergillus* genomics above, the genome sequencing of *A. nidulans*FGSC A4 was started by Cereon Genomics (Monsanto) in 1998 with the open discharge of the genome sequencing in June 2003. Two years later of explanation andstudyattempt, the genome paper by [21] was published. The *A. nidulans*genome was initially anticipated holdingnine thousand five hundred and forty one protein coding genes, yet following an impressive physical curation of the gene expectations in March of 2006, the amount of anticipated genes was changed to ten thousand seven hundred and one. A new article by [15] utilizes genomics and metabolic pathway reforms to furnish useful assignments to four hundred and seventy two stray genes.

Aspergillus Niger Genomics: A. Niger CBS 513.88 was the initial of so far three *A. Niger* strains to be a complete genome ordered. The genome ordering was started by the Dutch organization DSM in participation with Gene agreement in July, 2000, and was accounted for by DSM to be finalized in December of 2001. The strain is a forebear of the industrial glucoamylase- generating strains at present being used, and the genome order was broken down by [34]. The main input of this genome periodical was the explanation of several monetarily applicable enzymes, incorporating extra than two hundred accepted proteases and more pretty nearly one hundred and seventytheoretical and at one time distinguished carbohydrate-active-enzymes.

The second *A. niger*genome sequence is of *A. Niger* ATCC 9029, a frequently used laboratory strain. The genome was ordered by Integrated Genomics and published in 2005 as an intensely disjointed (9510 contigs) summary order without explanation or gene forecasts. At last, the Joint Genome Institute (JGI) dependent on the financing back from the United State Department of Energy (DOE) has finished and published the genome sequence of *A. niger* ATCC 1015, a citric acid maker initially depicted in 1917. The periodical of [11] was the initial research utilizing and introducing the *A. niger* ATCC 1015 genome sequence, despite the fact that not officially a publication the genome order.

Different genomics executed in *A. Niger* includes the ordering of twelve thousand eight hundred and twenty ESTs by [38] with the expectation of five thousand one hundred and eight genes. The study of the definite genes furnishes a significant substance that has significance both independently and in combination with an accessible genome sequence.

Besides, [26] ordered the total genome sequence of the 1a mtDNA sort of *A. Niger* and 2b mtDNA sort of *Aspergillus tubingensis* and discovered the two mtDNA to be very nearly indistinguishable.

The worth of a genome arrangement for an industrially pertinent species has been showed by DSM wherever it has been employed in manufactured goods development.

Illustrations incorporate a protease product known as "Brewer's Clarex" that averts cool vapor in beer [28], a protease product to wipe out unpleasantness of protein enhancement exercise beverages [19], and "Preventase", an asparaginase product intended at decreasing acryl amide development in baked and fried food [17].

Table 2: An Overview of Whole Genome Sequence Projects for *Aspergillus* Species [2]

Species	Strain	Sequencing	Contigs	Coverage (fold)	GenModel	Size (Mb)	Publication
A.aculeatus	ATCC 16872[a]	DOE Joint Genome Institute	N/A	N/A	N/A	N/A	Scott Baker. Personal commun.
A.Carbonarius	IMI 388653[a]	DOE Joint Genome Institute	N/A	N/A	N/A	N/A	Scott Baker. Personal commun.
A.davatus	NRRL 1	TIGR	231	114	9125	27.86	Fedorova et al. (2008)
A.(Neosartorya) Fischeri	NRR 181	TIGR	1067	N/A	10.407	32.55	Fedorova et al. (2008)
A.Flavus	NRR 3357	TIGR	3388	10	13.071	36.3	YU et al (2005)
A.Fumigatus	Af293	TIGR/Sanger instatute	19	10.5	9887	29.4	Nierman et al. (2005)
A.Fumigatus	A1163	TIGR/Celera Genomics/Merck	140	N/A	10.099	29.2	Fedorova et al (2008)

IJRS

		& co USA					
A.(Emericella) nidulans	FGSC A4	Broab institute	248	13	10.701	30.06	Galagan et al. (2005)
A. Niger	CBS 513.88	DSM, Neitherlands	498	7.5	14, 165	33.9	Peletan (2007)
A. Niger	ATCC 90929	Integrated Genomic	9510	4	N/A	N/A	
A. Niger	ATCC 1015	DOE Joint Genome instatute	24	8.9	11.200	34.9	Baker (2006)[b]
A. 0ryzae	RIB40	NTTE. japan	24	9	12.074	37	Machida et al.(2005
A. Parasiticus	unknown[a]	University of oklahoma	N/A	N/A	N/A	N/A	
A. terreus	ATCC 20542	Microbia	N/A	N/A	N/A	N/A	
A. terreus	NIH 2624	Broad institute	267	N/A	10.406	29.3	

IV. ASPERGILLUS GENOMIC RESOURCES

[23] Find out that no database holds each of the 14 available Aspergillus genomes; most are available from various resources. These includes the *Aspergillus* Genome Database (AspGD) reported by [3], FungiDB reported by [39], Central *Aspergillus* Data Repository (CADRE) reported by [29] and the *Aspergillus* Comparative Database (http://www. broadinstitute.org/annotation/genome/aspergillus_group/), with the remaining available from GenBank reported by [7] or the JGI genome portal reported by [24].

[5] Summarized some useful *Aspergillus* genome websites in table 3.

Table 3: Useful *Aspergillus* Genome Websites

SPECIES	Website URL Address
A. Fumigatus	http://www.tgr.org/tdb/e2kl/aful/ http://www.sanger.ac.uk/project/A fumigatus/
A. Nidulans	http://www.broad.mit.edu/annotation/fungi/aspergillus/
A. Niger	http://www.dsm.com/dfs/annovation/genomics/a http://www.intergratedgenomics.com/products.htmla
A. Oryzae	Whole genome not yet publicly available http://www.aist.go.jp/RIODB/ffdb/welcome.html
General *Aspergillus* sites	http://www.aspergillus.man.ac.uk/ http://www.fgs.net/aspergenome.html http://www.aspergellus-genomics.org/ http://www.genome.ou.edu/fungal.html http://www.cadre.man.ac.uk/

V. USES OF ASPERGILLUS GENOMICS DATA

A. fumigates is best generally linked of the *Aspergilli* with ailment among people despite the fact that different species of *Aspergillus* are likewise implicated [10]. Examination of the genome sequence of *A. fumigatus* might furnish signs to conceivable pathogenicity cause; however present researches indicate a multi-factorial origin of pathogenicity [1].

Publication of the annotated *A. fumigatus* genome, and analysis with different *Aspergillus* genomesare eminent and will give a detailed appraisal in genes that are anticipated to have a role in pathogenicity and allergenicity. Typical gene inactivation methodologies can then be utilized to experiment their imperativeness in pathogenicity. Gene knockouts could be challenging in *Aspergillus,* in spite of the fact that this could be enhanced by the production of huge flanks within the gene of importance [27]. Notwithstanding the study of the pathogenic mechanism of *A. fumigatus, Aspergillus spp.* might likewise give helpful frameworks for the investigation of human infection, as has been realized with *S. cerevisiae* in [6].

A. nidulans have been utilized a replica organism to investigate the molecular origin of human natural mistakes of metabolism in [37]. The possibility for this advance will be helped by the accessibility of post-genomic systems.

Aspergillus species are broadly used for the making of enzymes and metabolites, but genome information has just newly got available. In the differentiation, the yeast *S. cerevisiae* is less metabolically adaptable but its genome order has been available for a more extended time and the advancement of different apparatus has ease a mixed approach to useful genomics [13].

Genomics possibly will make available an insight into the molecular-genetic origin for species and strain discovery.

IJRS

Mutagenesis has routinely been the track for strain enhancement in fungal biotechnology and, as gene arrays start to be employed more broadly and will start to comprehend more the molecular origin for such upgrades [4].

VI. FUTURE APPLICATIONS OF ASPERGILLUS GENOMICS DATA

With the amount of *Aspergillus species*yet in the sequencing channel, it is understandable that the genomic period of *Aspergilli* has just barely begun. The upcoming discharge of additional sequences and in addition more exhaustive relative genomics researches holds guarantees for sorting out numerous diverse novel mechanisms and recognition of basics that characterizes one species and are preserved throughout the genus. Particularly for study in pathogenicity, eukaryotic cell physiology, and biotechnology, upcoming genome research might have a considerable effect [2]. The arrangements of *Aspergillus* genomes, together the right now available and the ones to be released sometime later will proceed to make the advancement of alternate levels of system-wide study.

VII. CONCLUSION

The genus *Aspergillus* described a different assembly of fungi that are within the most plentiful fungi on the planet. The accomplishment of *Aspergilli* is clarified, by the way, that they are not extremely particular concerning their a biotic development conditions that they can break a wide range of organic molecules , and by the way that they make high numbers of asexual and sexual spores that are scattered over small and long distances [26]

Not exactly a decade after the release of the first freely available *Aspergillus* genomes, the investigation of the role and development of the genome of this extensively significant genus has recently started producing exceptional novel experiences into genome architecture, sexual reproduction, populace biology, and secondary metabolism and development, and virulence mechanisms [23].

The developing collaboration between the generous and always unfolding forms of learning on *Aspergillus* genomics, systematic, natural history, molecular genetics and development, natural products chemistry, human, animal and plant ailment, and biotechnology, coupled with the striking phenotypic adaptability exhibit in the genus, make it perfect for tending to central inquiries over some levels of biological organization [13].

REFERENCES

[1] Anderson M.J., Brookman J.L., Denning D.W. (2003). Aspergillus. In Prade R.A., Bohnert H.J. (ed) Genomics of Plants and Fungi. New York. Marcel Dekker Inc. Publishers pp 1-39.

[2] Andersen M.R., Nielsen J. (2009). Current status of systems biology in Aspergilli.Fungal Genetics and Biology, **46** (2009) pp180-190

[3] Arnaud, M.B., (2012) The Aspergillus Genome Database (AspGD): recent developments in comprehensive multispecies curation, comparative genomics and community resources. Nucleic Acids Research **40**pp D653–D659

[4] Archer, D.B., Dyer, P.S., (2004). From genomics to post-genomics in Aspergillus.Current

[5] Opinion in Microbiology **7**, pp 499–504.

[6] Archer D.B., Turner G.,(2006).Genomics of protein secretion and Hyphal Growth in Aspergillus.Fungal Genomics, pp 75-96

[7] Bassett D.E., Boguski M.S., Hieter P. (1996) Yeast genes and human disease. Nature, **379**pp 589-590.

[8] Benson, D.A., (2009) GenBank. Nucleic Acids Research **37**pp D26–D31

[9] Bennett J.W.,(2010). An overview of the Genus Aspergillus.In Machida M., Gomi K. (ed.) Aspergillus Molecular Biology and Genomics England.Caister, pp 1-17

[10] Borkovich K.A., Alex L.A., Yarden O., Freitag M., Turner G.E., Read N.D.,Seiler S., Bell-Pedersen D., Paietta J., Plesofsky N. (2004). Lessons from the genome sequence of Neurospora crassa: tracing the path from genomic blueprint to a multicellular organism.

[11] Microbiology and Molecular Biology Review, **68**pp 1-108.

[12] Brakhage A.A., Langfelder K. (2002). Menacing mold: the molecular biology of Aspergillus fumigatus.Annual Review of Microbiology, **56**pp 433-455.

[13] Baker, S., (2006). Aspergillus niger genomics: past, present and into the future. Medical Mycology **44** (1), pp S17–S21.

[14] Bowyer, P., Denning, D., (2007). Genomic analysis of allergen genes in Aspergillus

[15] spp: the relevance of genomics to everyday research. Medical Mycology **45** (1), pp 17–

[16] 26.

[17] Castrillo J.O, Oliver S.G. (2004). Yeast as a touchstone in post-genomic research: strategies for integrative analysis in functional genomics.Journal of Biochemistry and Molecular Biology, **37**pp 93-106.

[18] Dujon, B. (2010) Yeast evolutionary genomics. Nature Review Genetics **11**pp 512–524

[19] David, H., Ozcelik, I., Hofmann, G., Nielsen, J., (2008). Analysis of Aspergillus nidulans

[20] metabolism at the genome-scale. BMC Genomics**9** (1), pp 163.

[21] Denning D.W., Anderson M.J., Turner G., Latgé, J. P., Bennett J.W., (2002).

[22] Sequencing the Aspergillus fumigatus genome.The Lancet Infectious Disease**2**, pp 251–253.

[23] DSM Food Specialities (2007). First ‘‘acrylamide-free” biscuits will be

[24] launched in Germany. Press release http://www.dsm.com/en_US/html/dfs/

[25] news_items/acrylamide_free_biscuits.htm.

IJRS

[26] Dyer P.S., Paoletti M., Archer, D.B., (2003). Genomics reveals sexual secrets of Aspergillus. Microbiology Comment **149**, pp 2301–2303.

[27] Edens L., Dekker P., van der Hoeven R., Deen F., De Roos A., Floris R., (2005).

[28] Extracellular prolylendoprotease from Aspergillus nigerand its use in the debittering of protein hydrolysates. Journal of Agriculture and Food Chemistry.**53**pp 7950–7957.

[29] Fedorova N., Khaldi N., Joardar V., Maiti R., Amedeo P., Anderson M., Crabtree J.,Silva J., Badger J., Albarraq A., Angiuoli S., Bussey H., Bowyer P., Cotty P.,Dyer P., Egan A., Galens K., Fraser-Liggett C., Haas B., Inman J., Kent R.,Lemieux S., Malavazi I., Orvis J., Roemer T., Ronning C., Sundaram J., SuttonG., Turner G., Venter J., White O., Whitty B., Youngman P., Wolfe K., GoldmanG., Wortman J., Jiang B., Denning D., Nierman W.C., (2008). Genomic islands inthe pathogenic filamentous fungus Aspergillus fumigatus. PLoS Genetics**4** (4),e1000046.

[30] Galaga, J., Calvo, S., Cuomo, C., Ma, L.-J., Wortman, J., Batzoglou, S., Lee, S.-I.,Basturkmen, M., Spevak, C., Clutterbuck, J., Kapitonov, V., Jurka, J., Scazzocchio,C., Farman, M., Butler, J., Purcell, S., Harris, S., Braus, G., Draht, O., Busch, S.,D'Enfert, C., Bouchier, C., Goldman, G., Bell-Pedersen, D., Griffiths-Jones, S.,Doonan, J., Yu, J., Vienken, K., Pain, A., Freitag, M., Selker, E., Archer, D., Penalva,M., Oakley, B., Momany, M., Tanaka, T., Kumagai, T., Asai, K., Machida, M.,Nierman, W., Denning, D., Caddick, M., Hynes, M., Paoletti, M., Fischer, R., Miller,B., Dyer, P., Sachs, M., Osmani, S., Birren, B., (2005). Sequencing of Aspergillusnidulans and comparative analysis with A. fumigatus and A. oryzae.Nature **438**(7071) pp 1105–1115.

[31] Galaga J.E., Calvo S.E., Borkovich K.A., (2003). The genome sequence of thefilamentous fungus Neurospora crassa.Nature **422**pp 859–868.

[32] Gibbsons J.G., Rokas A., (2013). The function and evolution of the Aspergillus genome.Trends in Microbiology, **21** (1) pp 14-21

[33] Grigoriev I.V. (2012) The genome portal of the Department of Energy Joint Genome Institute. Nucleic Acids Research**40** pp D26–D32

[34] Juhász, A., Pfeiffer, I., Keszthelyi, A., Kucsera, J., Vágvölgyi, C., Hamari, Z., (2008).Comparative analysis of the complete mitochondrial genomes of AspergillusnigermtDNA type 1a and Aspergillus tubingensismtDNA type 2b. FEMSMicrobiology Letter**281** (1) pp 51–57.

[35] Krijgsheld P., Bleichradt G.T.,Veluw V., Wang F., Muller J., Dijksterhuis and Wosten H.A.B., (2012). Development in Aspergillus.Studies in Mycology**74** pp 1-29

[36] Langfelder K, Gattung S, Brakhage AA. (2002). A novel method used to delete a new Aspergillus fumigatus ABC transporter-encoding gene. Current Genetics **41**pp 268-274.

[37] Lopez, M., Edens, L., (2005). Effective prevention of chill-haze in beer using an acid proline specific endoprotease from Aspergillus niger. J. Agric. Food Chem.**53**, PP 7944–7949.

[38] MabeyGilsenan, J. (2012) CADRE: the Central Aspergillus Data Repository 2012. Nucleic Acids Research**40** pp D660–D666

[39] Machida, M., Asai, K., Sano, M., Tanaka, T., Kumagai, T., Terai, G., Kusumoto, K.-I.,Arima, T., Akita, O., Kashiwagi, Y., Abe, K., Gomi, K., Horiuchi, H., Kitamoto, K.,Kobayashi, T., Takeuchi, M., Denning, D., Galagan, J., Nierman, W., Yu, J.,Archer, D., Bennett, J., Bhatnagar, D., Cleveland, T., Fedorova, N., Gotoh, O.,Horikawa, H., Hosoyama, A., Ichinomiya, M., Igarashi, R., Iwashita, K., Juvvadi,P., Kato, M., Kato, Y., Kin, T., Kokubun, A., Maeda, H., Maeyama, N.,Maruyama, J.-I., Nagasaki, H., Nakajima, T., Oda, K., Okada, K., Paulsen, I.,Sakamoto, K., Sawano, T., Takahashi, M., Takase, K., Terabayashi, Y., Wortman,J., Yamada, O., Yamagata, Y., Anazawa, H., Hata, Y., Koide, Y., Komori, T.,Koyama, Y., Minetoki, T., Suharnan, S., Tanaka, A., Isono, K., Kuhara, S.,Ogasawara, Kikuchi, H., (2005). Genome sequencing and analysis of Aspergillusoryzae.Nature**438** (7071) pp 1157–1161.

[40] Nierman, W., Pain, A., Anderson, M., Wortman, J., Kim, H., Arroyo, J., Berriman, M.,Abe, K., Archer, D., Bermejo, C., Bennett, J., Bowyer, P., Chen, D., Collins, M.,Coulsen, R., Davies, R., Dyer, P.S., Farman, M., Fedorova, N., Fedorova, N.,Feldblyum, T., Fischer, R., Fosker, N., Fraser, A., Garcia, J., Garcia, M., Goble, A.,Goldman, G., Gomi, K., Griffith-Jones, S., Gwilliam, R., Haas, B., Haas, H., Harris,D., Horiuchi, H., Huang, J., Humphray, S., Jimenez, J., Keller, N., Khouri, H.,Kitamoto, K., Kobayashi, T., Konzack, S., Kulkarni, R., Kumagai, T., Lafton, A.,Latge, J.-P., Li, W., Lord, A., Lu, C., Majoros, W., May, G., Miller, B., Mohamoud, Y.,Molina, M., Monod, M., Mouyna, I., Mulligan, S., Murphy, L., O'Neil, S., Paulsen, I.,Penalva, M., Pertea, M., Price, C., Pritchard, B., Quail, M., Rabbinowitsch, E.,Rawlins, N., Rajandream, M.-A., Reichard, U., Renauld, H., Robson, G., deCordoba, S., Rodriguez-Pena, J., Ronning, C., Rutter, S., Salzberg, S., Sanchez, M.,Sanchez-Ferrero, J., Saunders, D., Seeger, K., Squares, R., Squares, S., Takeuchi, M.,Tekaia, F., Turner, G., de Aldana, C., Weidman, J., White, O., Woodward, J., Yu, J.-H., Fraser, C., Galagan, J., Asai, K., Machida, M., Hall, N., Barrell, B., Denning, D.,(2005). Genomic sequence of the pathogenic and allergenic filamentous fungusAspergillus fumigatus.Nature **438** (7071) pp 1151–1156.

[41] Pain A, Woodward J, Quail MA, Anderson MJ, Clark R, Collins M, Fosker N, Fraser A, Harris D, Larke N., (2004) Insight into the genome of Aspergillus fumigatus: analysis of a 922 kb region encompassing the nitrate assimilation gene cluster. Fungal Genet Biol ,**41**pp443-453.

[42] Paoletti, M., Rydholm, C., Schwier, E., Anderson, M., Szakacs, G., Lutzoni, F.,Debeaupuis, J., Latgé, J., Denning, D., Dyer, P., (2005). Evidence for sexuality inthe opportunistic fungal pathogen Aspergillus fumigatus.Current Biology**15** (13) pp1242–1248.

[43] Pel, H., de Winde, J., Archer, D., Dyer, P., Hofmann, G., Schaap, P., Turner, G., de Vries,R., Albang, R., Albermann, K., Andersen, M., Bendtsen, J., Benen, J., van den Berg,M., Breestraat, S., Caddick, M., Contreras, R., Cornell, M., Coutinho, P., Danchin,E., Debets, A., Dekker,

IJRS

P., van Dijck, P., van Dijk, A., Dijkhuizen, L., Driessen, A.,d'Enfert, C., Geysens, S., Goosen, C., Groot, G., de Groot, P., Guillemette, T.,Henrissat, B., Herweijer, M., van den Hombergh, J., van den Hondel, C., van derHeijden, R., van der Kaaij, R., Klis, F., Kools, H., Kubicek, C., van Kuyk, P., Lauber,J., Lu, X., van der Maarel, M., Meulenberg, R., Menke, H., Mortimer, M., Nielsen, J.,Oliver, S., Olsthoorn, M., Pal, K., van Peij, N., Ram, A., Rinas, U., Roubos, J., Sagt, C.,Schmoll, M., Sun, J., Ussery, D., Varga, J., Vervecken, W., van de Vondervoort, P.,Wedler, H., Wosten, H.A., Zeng, A., van Ooyen, A., Visser, J., Stam, H., (2007).Genome sequencing and analysis of the versatile cell factory Aspergillus nigerCBS 513.88. Nature Biotechnology**25** (2) pp 221–231.

[44] Rokas, A., Payne, G., Fedorova, N., Baker, S., Machida, M., Yu, J., Georgianna, D.R.,Dean, R.A., Bhatnagar, D., Cleveland, T., Wortman, J., Maiti, R., Joardar, V.,Amedeo, P., Denning, D., Nierman, W., (2007). What can comparative genomicstell us about species concepts in the genus? Studies in Mycology.**59** (1) pp 11–17.

[45] Ronning, C.M., Fedorova, N., Bowyer, P., Coulson, R., Goldman, G., Kim, H., Turner, G.,Wortman, J., Yu, J., Anderson, M., Denning, D., Nierman, W., (2005). Genomics ofAspergillus fumigatus.Rev. Iberoam. Micol**22**pp 223–228.

[46] Rodriguez J.M, Ruiz-Sala P, Ugarte M, Pen˜ alva M.A. (2004) Fungal metabolic model for 3-methylcrotonyl-CoA carboxylase deficiency. J BiolChem,**279**pp 4578-4587.

[47] Semova, N., Storms, R., John, T., Gaudet, P., Ulycznyj, P., Min, X., Sun, J., Butler, G.,Tsang, A., (2006). Generation, annotation, and analysis of an extensive AspergillusnigerEST collection. BMC Microbiology**2** (6) pp 7.Stajich, J.E. et al. (2012) FungiDB: an integrated functional genomics database for fungi. Nucleic Acids Research **40**pp D675–D681

[48] Stajich, J.E. (2012) FungiDB: an integrated functional genomics database for fungi. Nucleic Acids Res.**40**pp D675–D681

[49] Wortman, J.R., Fedorova, N., Crabtree, J., Joardar, V., Maiti, R., Haas, B.J., Amedeo, P.,Lee, E., Angiuoli, S.V., et al., (2006). Whole genome comparison of the A. fumigatesfamily. Medical Mycology**44** pp S3–S37.

[50] Yu, J., Cleveland, T., Nierman, W., Bennett, J., (2005). Aspergillus flavus genomics:gateway to human and animal health, food safety, and crop resistance todiseases. Rev. Iberoam. Micoiogy**22** (4) pp 192–202.

IJRS